

Bounded nonvanishing functions and Bateman functions

WOLFRAM KOEPF, DIETER SCHMERSAU

to appear in: Complex Variables, 1994

Abstract:

We consider the family $\tilde{B}$ of bounded nonvanishing analytic functions $f(z) = a_0 + a_1 z + a_2 z^2 + \dots$ in the unit disk. The coefficient problem had been extensively investigated (see e. g. [2], [13], [14], [16], [17], [18], [20]), and it is known that

$$|a_n| \leq \frac{2}{e}$$

for $n = 1, 2, 3$, and 4. That this inequality may hold for $n \in \mathbb{N}$, is known as the Krzyż conjecture.

It turns out that for $f \in \tilde{B}$ with $a_0 = e^{-t}$

$$f(z) \prec e^{-t \frac{1+z}{1-z}}$$

so that the superordinate functions $e^{-t \frac{1+z}{1-z}} = \sum_{k=0}^{\infty} F_k(t) z^k$ are of special interest. The corresponding coefficient functions $F_k(t)$ had been independently considered by Bateman [3] who had introduced them with the aid of the integral representation

$$F_k(t) = (-1)^k \frac{2}{\pi} \int_0^{\pi/2} \cos(t \tan \theta - 2k\theta) d\theta .$$

We study the Bateman functions and formulate properties that give insight in the coefficient problem in $\tilde{B}$.

1 Introduction

We consider functions that are analytic in the unit disk

$$\mathbb{D} := \{z \in \mathbb{C} \mid |z| < 1\} .$$

An analytic function f is called subordinate to g , if $f = g \circ \omega$ for some analytic function ω with $\omega(0) = 0$ and $\omega(\mathbb{D}) \subset \mathbb{D}$; we write $f \prec g$. The subordination principle states that if g is univalent then $f \prec g$ if and only if $f(0) = g(0)$ and $f(\mathbb{D}) \subset g(\mathbb{D})$, see e. g. [15], § 23.

Let $\tilde{B}$ denote the family of bounded nonvanishing analytic functions $f(z) = a_0 + a_1 z + a_2 z^2 + \dots$ in $\mathbb{D}$. As f is nonvanishing, we have $\operatorname{Re} \ln f(z) < 0$, and by the subordination principle it turns out that for $a_0 = e^{-t}$

$$-\ln f(z) \prec t \frac{1+z}{1-z} ,$$

and so $f(z) = e^{-t} + a_1 z + a_2 z^2 + \cdots \in \tilde{B}$ if and only if

$$f(z) \prec e^{-t} \frac{1+z}{1-z} . \quad (1)$$

Thus the superordinate functions

$$G(t, z) = e^{-t} \frac{1+z}{1-z} =: \sum_{k=0}^{\infty} F_k(t) z^k \quad (2)$$

are of special interest. Graphs of the functions $F_n(t)$ ($n = 1, \dots, 5$) are given in Figure 1.




Figure 1: Graphs of the functions $F_n(t)$ ($n = 1, \dots, 5$)

The following is a list of the first functions F_n :

$$\begin{aligned} F_0(t) &= e^{-t} \\ F_1(t) &= -2te^{-t} , \\ F_2(t) &= 2e^{-t}(-1+t)t , \\ F_3(t) &= \frac{2te^{-t}(-3+6t-2t^2)}{3} , \\ F_4(t) &= \frac{2te^{-t}(-3+9t-6t^2+t^3)}{3} , \\ F_5(t) &= \frac{2te^{-t}(-15+60t-60t^2+20t^3-2t^4)}{15} . \end{aligned}$$

We consider the coefficient problem ($n \in \mathbb{N}$) to find

$$A_n := \max_{f \in \tilde{B}} |a_n(f)| .$$

That the maximum exists for all $n \in \mathbb{N}$ follows from the fact that the union of $\tilde{B}$ with the constant functions $c \in \overline{\mathbb{D}}$ forms a compact family of analytic functions. For the coefficient problem it is no

loss of generality to assume that $a_0 > 0$ so that we can assume that (1) holds for some $t > 0$. For small n it is then easy to solve the coefficient problem using subordination techniques.

As $f \prec g$ implies that $|a_1(f)| \leq |a_1(g)|$ (see e. g. [15], Theorem 212), we have

$$A_1 = \max_{t \geq 0} \left| a_1 \left(e^{-t \frac{1+z}{1-z}} \right) \right| = \max_{t \geq 0} |F_1(t)| = \max_{t \geq 0} 2t e^{-t} = \frac{2}{e}$$

with equality iff $t = 1$, and $f(z) = \eta e^{-\frac{1+\xi z}{1-\xi z}}$ ($|\xi| = |\eta| = 1$).

By the composition with a Möbius transform, this leads to the inequality (see [16])

$$(1 - |z|^2) |f'(z)| \leq \frac{2}{e} \quad (z \in \mathbb{D}) \quad (3)$$

from which we may deduce by a standard technique (see e. g. [7], p. 72, Exercise 17) that

$$\begin{aligned} n |a_n(f)| &= |a_{n-1}(f')| = \left| \frac{1}{2\pi i} \int_{\partial \mathbb{D}_r} \frac{f'(\zeta)}{\zeta^n} d\zeta \right| \leq \frac{1}{r^{n-1}} \left(\frac{1}{2\pi} \int_0^{2\pi} |f'(r e^{i\theta})| d\theta \right) \\ &\leq \frac{1}{r^{n-1}(1-r^2)} \frac{2}{e} \leq \frac{2}{e} \frac{n+1}{2} \left(1 + \frac{2}{n-1} \right)^{\frac{n-1}{2}} < n \end{aligned}$$

where we used (3) and chose $r^2 = \frac{n-1}{n+1}$. Unfortunately this estimate is just too weak to be off value: The bound 1 for A_n is very elementary, and holds even for all functions bounded by 1ff. Each global bound less than 1 would be new, however.

It is similarly easy to solve the coefficient problem for $n = 2$ using subordination techniques, ([15], Theorem 212, see e. g. [10]).

Using several methods it was shown that

$$A_n = \frac{2}{e} \quad (4)$$

for $n = 1, 2, 3$, and 4. Obviously $G(1, z^n)$ has n^{th} coefficient equal to $\frac{2}{e}$, which makes these results sharp. That (4) may hold for $n \in \mathbb{N}$, is known as the Krzyż conjecture.

If the subordinate function has very regular coefficient behavior, then global coefficient results are available: If

$$\sum_{k=0}^{\infty} a_k z^k \prec \sum_{k=0}^{\infty} b_k z^k, \quad (5)$$

and if the coefficient sequence b_n is nonnegative, nonincreasing, and convex, then $|a_n| \leq b_0$ for all $n \in \mathbb{N}_0$, and if the coefficient sequence b_n is nonnegative, nondecreasing, and convex, then $|a_n| \leq b_n$ all $n \in \mathbb{N}_0$ (see e.g. [15], Theorem 216). On the other hand, the coefficient sequences of our subordinate functions $G(t, z)$ are highly irregular for all $t > 0$.

Another important result, however, can be obtained by subordination techniques, as well. It is well known that if (5) holds, then

$$\sum_{k=0}^{\infty} |a_k|^2 \leq \sum_{k=0}^{\infty} |b_k|^2.$$

Especially: If an analytic function $f(z) = \sum_{k=0}^{\infty} a_k z^k$ of the unit disk is bounded by 1, then $f \prec z$, and the relation

$$\sum_{k=0}^{\infty} |a_k|^2 \leq 1$$

(following also directly from Parseval's identity) is obtained (for sharper versions see also [7], Theorem 6.1 and Theorem 6.2). Equality holds if and only if (see [19], Theorem 3) f is an inner function, i.e. if the radial limit $f(e^{i\theta}) := \lim_{r \rightarrow 1} f(re^{i\theta}) = 1$ for almost all $e^{i\theta}$ on the unit circle $\partial\mathbb{D}$. Nonvanishing inner functions with positive $f(0)$ have the representation (see e.g. [9], second theorem on p. 66)

$$f(z) = \exp \left(- \int_{\partial\mathbb{D}} \frac{1 + e^{i\theta} z}{1 - e^{i\theta} z} d\mu(\theta) \right) ,$$

where μ is a singular positive measure on the unit circle $\partial\mathbb{D}$. If we choose a point measure μ concentrating its full measure t at the point $\theta = 1$, we get the function $G(t, z) = e^{-t \frac{1+z}{1-z}}$ of Equation (2) so that we are lead to the identity

$$\sum_{k=0}^{\infty} F_k^2(t) = 1 .$$

For each individual coefficient of $G(t, z)$ we thus have the (weak) inequality

$$|F_n(t)| \leq \sqrt{1 - F_0^2(t)} = \sqrt{1 - e^{-2t}} \quad (n \in \mathbb{N}) . \quad (6)$$

It is the purpose of this paper to develop further properties, especially inequalities, for the functions F_n ($n \in \mathbb{N}$), giving more insight in the coefficient problem for $\tilde{B}$.

2 A collection of properties of the Bateman functions

In [3] (see also [1], § 13.6) Bateman introduced the functions ($x \geq 0$)

$$k_n(x) := \frac{2}{\pi} \int_0^{\pi/2} \cos(x \tan \theta - n \theta) d\theta ,$$

and he verified that ([3], formula (2.7))

$$k_{2m}(x) = (-1)^m e^{-x} (L_m(2x) - L_{m-1}(2x)) \quad (7)$$

where $L_m(t)$ denotes the m^{th} Laguerre polynomial. On the other hand if one defines the functions F_n ($n \in \mathbb{N}_0$) with the aid of the generating function

$$e^{-t \frac{1+z}{1-z}} =: \sum_{k=0}^{\infty} F_k(t) z^k ,$$

one gets immediately (see [10], formula (14), and p. 178)

$$F_n(t) = e^{-t} \left(L_n(2t) - L_{n-1}(2t) \right) \quad (8)$$

and a comparison of (7) and (8) yields the relation

$$F_n(t) = (-1)^n k_{2n}(t)$$

so that we get the Bateman representation

$$F_n(t) = (-1)^n \frac{2}{\pi} \int_0^{\pi/2} \cos(t \tan \theta - 2n\theta) d\theta \quad (9)$$

for our functions F_n . By Bateman's work we are prepared to state many further properties: For $n \in \mathbb{N}$ the function F_n satisfies the differential equation (see [3], formula (5.1))

$$t F_n''(t) = (t - 2n) F_n(t) \quad (10)$$

with the initial values

$$F_n(0) = 0 \quad \text{and} \quad F_n'(0) = -2, \quad (11)$$

and the Rodriguez type formula (see [3], formula (31))

$$F_n(t) = \frac{t e^t}{n!} \frac{d^n}{dt^n} \left(e^{-2t} t^{n-1} \right).$$

The differential equation can also be obtained completely algorithmically (see [11]–[12]).

Further we get the following connection with the generalized Laguerre polynomials (see [23], ffp. 216, formula (1.15))

$$F_n(t) = e^{-t} L_n^{(-1)}(2t), \quad (12)$$

and (see [22], formula (5.2.1))

$$F_n(t) = -e^{-t} \frac{2t}{n} L_{n-1}^{(1)}(2t), \quad (13)$$

from which one may deduce the hypergeometric representation

$$F_n(t) = -2t e^{-t} {}_1F_1 \left(\begin{matrix} 1 - n \\ 2 \end{matrix} \middle| 2t \right),$$

and the explicit representation

$$F_n(t) = \frac{e^{-t}}{n} \sum_{k=1}^n \frac{(-1)^k}{(k-1)!} \binom{n}{k} (2t)^k.$$

Bateman obtained further relations: a difference equation ([3], formula (4.1))

$$(n-1)(F_n(t) - F_{n-1}(t)) + (n+1)(F_n(t) - F_{n+1}(t)) = 2t F_n(t) \quad (14)$$

that is also an easy consequence of the defining equation using the generating function, he obtained a difference differential equation ([3], formula (4.2))

$$(n+1)F_{n+1}(t) - (n-1)F_{n-1}(t) = 2tF'_n(t) ,$$

and a system of differential equations ([3], formula (4.3))

$$F'_n(t) - F'_{n+1}(t) = F_n(t) + F_{n+1}(t) , \quad (15)$$

from which he is lead to the inequalities for F_n ([3], formula (4.4))

$$|F_n(t)| \leq \frac{2n}{t} \quad (n > 2) , \quad (16)$$

and for F'_n ([3], formula (4.5))

$$|F'_n(t)| \leq \frac{n}{t} \quad (n > 2) . \quad (17)$$

For large t the first inequality is a refinement of the trivial estimate

$$|F_n(t)| \leq 1 \quad (18)$$

that follows from (6) or from the Bateman representation (9).

Finally Bateman obtained the following statements about Integrals of products $F_n F_m$ ($n, m \in \mathbb{N}$) (see [3], formula (2.91))

$$\int_0^\infty F_n^2(t) dt = 1 \quad \text{and} \quad \int_0^\infty F_n(t) F_m(t) dt = \begin{cases} 0 & \text{if } |n-m| \neq 1 \\ \frac{1}{2} & \text{otherwise} \end{cases} . \quad (19)$$

We state further properties: The functions F_n ($n \in \mathbb{N}$) have a zero at the origin and $n-1$ further positive real zeros (see e. g. [23], *Nullstellensatz*, p. 123) (indeed, by (13), $F_n(t)$ has the same zeros as $L_{n-1}^{(1)}(2t)$).

From the differential equation (10) we moreover see that at $t = 2n$ there is a point of inflection, and as $F_n(t) \rightarrow 0$ for $t \rightarrow \infty$, and all other points of inflection lie at the zeros of F_n one easily deduces that $t = 2n$ must be the largest point of inflection of F_n implying that all the zeros of F_n lie in the interval $[0, 2n)$. The successive relative maxima of $|F_n|$ lying between the zeros of F_n form an increasing sequence (see [22], Theorem 7.6.2, $\alpha = -1$), so that the largest value attained by $|F_n(t)|$ is attained at the last zero of F'_n which is seen to lie between the last zero T_n of F_n and the point $t = 2n$. For small n the mentioned qualitative properties of F_n can be recognized in Figure 1.

By a result of Hahn ([8], formula (17)) the last zero T_n (being the last zero of $L_{n-1}^{(1)}(2t)$) satisfies the relation

$$4n - 2 - C_1 \sqrt[3]{4n-2} < 2T_n < 4n - 2 - C_2 \sqrt[3]{4n-2} \quad (20)$$

with two positive constants $C_1, C_2 \in \mathbb{R}^+$ that are independent of n , in particular

$$\lim_{n \rightarrow \infty} \frac{T_n}{n} = 2 . \quad (21)$$

The right hand side of (20) leads to the sharpened inequality

$$T_n < 2n - 1 ,$$

and Puiseux series expansion of (20) yields the refinement of (21)

$$\frac{T_n}{n} = 2 - O\left(\left(\frac{1}{n}\right)^{2/3}\right) .$$

3 Representation by residues

To the system of differential equations given by (15) together with the initial conditions $F_n(0) = 0$ ($n \in \mathbb{N}$) the technique of Laplace transformation

$$\mathcal{L}(f)(z) := \int_0^{\infty} e^{-zt} f(t) dt$$

can be applied to deduce a representation by residues for F_n . It is well-known that $\mathcal{L}(f') = z\mathcal{L}(f) - f(0)$ (see e.g. [6], Satz 9.1) so that we obtain ($n \in \mathbb{N}$)

$$(z+1)\mathcal{L}(F_{n+1}) = (z-1)\mathcal{L}(F_n)$$

or

$$\mathcal{L}(F_{n+1}) = \frac{z-1}{z+1} \mathcal{L}(F_n) .$$

Induction shows then that for $n \in \mathbb{N}$ and $k \in \mathbb{N}_0$

$$\mathcal{L}(F_{n+k}) = \left(\frac{z-1}{z+1} \right)^k \mathcal{L}(F_n) . \quad (22)$$

To obtain the initial function $\mathcal{L}(F_1)$, we use $F_0(t) = e^{-t}$ to get first

$$\mathcal{L}(F_0)(z) = \int_0^{\infty} e^{(z-1)t} dt = \frac{1}{1+z} .$$

Further from (15) with $n = 0$ we are lead to

$$(z+1)\mathcal{L}(F_1) = (z-1)\mathcal{L}(F_0) - 1$$

or

$$\mathcal{L}(F_1)(z) = -\frac{2}{(1+z)^2} .$$

Thus by an application of (22) with $n = 1$ we have finally

$$\mathcal{L}(F_k)(z) = -\frac{2}{(1+z)^2} \left(\frac{z-1}{z+1} \right)^{k-1} .$$

If we use now the inverse Laplace transform (see e.g. [6], p. 170, formula (15)), we get

$$F_k(t) = \lim_{R \rightarrow \infty} \frac{1}{2\pi i} \int_{\gamma_R} e^{tz} \mathcal{L}(F_k)(z) dz ,$$

where $\gamma_R : [-R, R] \rightarrow \mathbb{C}$ is given by $\gamma_R(\tau) = i\tau$, and therefore we have the integral representation

$$F_k(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} e^{it\tau} \frac{(\tau+i)^{k-1}}{(\tau-i)^{k+1}} d\tau .$$

By a standard procedure this can be identified as the residue (see e.g. [4ff], p. 217, formula (12))

$$\int_{-\infty}^{\infty} e^{it\tau} \frac{(\tau+i)^{k-1}}{(\tau-i)^{k+1}} d\tau = 2\pi i \operatorname{Res} \left(e^{itz} \frac{(z+i)^{k-1}}{(z-i)^{k+1}} \right) ,$$

and therefore we have the representation ($k \in \mathbb{N}$)

$$F_k(t) = 2i \operatorname{Res} \left(e^{itz} \frac{(z+i)^{k-1}}{(z-i)^{k+1}} \right) .$$

These results are collected in

Theorem 1 The Bateman functions F_k ($k \in \mathbb{N}$) satisfy the integral representation

$$F_k(t) = \frac{1}{\pi} \int_{-\infty}^{\infty} e^{it\tau} \frac{(\tau+i)^{k-1}}{(\tau-i)^{k+1}} d\tau$$

and therefore the residual representation

$$F_k(t) = 2i \operatorname{Res} \left(e^{itz} \frac{(z+i)^{k-1}}{(z-i)^{k+1}} \right) .$$

4 Results deduced from the differential equation

In this section we deduce another statements about an integral involving the Bateman functions and get an estimate for $|F_n|$ using its differential equation (10). Multiplying (10) by $2 F'_n(t)/t$, we have

$$2 F'_n(t) F''_n(t) = 2 F_n(t) F'_n(t) - \frac{2n}{t} 2 F_n(t) F'_n(t) .$$

We integrate from 0 to t , and get for $n \in \mathbb{N}$ using the initial values (11)

$$(F'_n)^2(t) - 4 = F_n^2(t) - \int_0^t \frac{2n}{\tau} 2 F_n(\tau) F'_n(\tau) d\tau . \quad (23)$$

For the last integral we get integrating by parts

$$\begin{aligned} \int_0^t \frac{2n}{\tau} 2 F_n(\tau) F'_n(\tau) d\tau &= \left. \frac{2n}{\tau} F_n^2(\tau) \right|_0^t + \int_0^t \frac{2n}{\tau^2} F_n^2(\tau) d\tau \\ &= \frac{2n}{t} F_n^2(t) - 2n F_n(0) F'_n(0) + \int_0^t 2n \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau \\ &= \frac{2n}{t} F_n^2(t) + 2n \int_0^t \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau . \end{aligned}$$

So we have the identity

$$(F'_n)^2(t) - 4 = F_n^2(t) - \frac{2n}{t} F_n^2(t) - 2n \int_0^t \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau. \quad (24)$$

From this identity by letting $t \rightarrow \infty$ we are lead to the statement

$$\int_0^\infty \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau = \frac{2}{n}$$

as $\lim_{t \rightarrow \infty} F_n(t) = \lim_{t \rightarrow \infty} F'_n(t) = 0$. Therefore in particular ($t \geq 0$)

$$\int_0^t \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau < \frac{2}{n}. \quad (25)$$

At this point we like to mention that from (24) it is now very easy to deduce the inequality $t < 2n$ for a local extremum of F_n , again (compare § 2), as an application of (25) yields

$$F_n^2(t) - \frac{2n}{t} F_n^2(t) = (F'_n)^2(t) - 4 + 2n \int_0^t \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau < (F'_n)^2(t),$$

and therefore for any point with $F'_n(t) = 0$ we get $t < 2n$.




Figure 2: The estimates (6) and (26) for $n = 1, \dots, 5$

To deduce an estimate for $|F_n|$ we regroup (24) and get

$$\frac{2n-t}{t} F_n^2(t) = 4 - (F'_n)^2(t) - 2n \int_0^t \left(\frac{F_n(\tau)}{\tau} \right)^2 d\tau < 4,$$

and for $t < 2n$ (which is the critical region) finally

$$|F_n^2(t)| < \frac{4t}{2n-t} . \quad (26)$$

We note that, however, this improves (6) for small t only, see Figure 2.

In the next section we will give a further improvement of (26).

5 Estimates by the Szegő method

We consider the generalized Laguerre polynomials $L_n^{(\alpha)}(x)$ ($\alpha \in \mathbb{R}$) given by their Rodriguez formula

$$x^\alpha e^{-x} L_n^{(\alpha)}(x) = \frac{1}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}) \quad (27)$$

(see [23], p. 213, formula (1.3)). Szegő ([21], see [22], p. 159, formula (7.21.3)) considered the case $\alpha = 0$, and was lead to the inequality

$$e^{-\frac{x}{2}} \left| L_n^{(0)}(x) \right| < 1 \quad (x > 0) . \quad (28)$$

Using a similar method we get the following development. For $n \in \mathbb{N}_0$ and $\alpha \in \mathbb{Z}$ the function $f_{n\alpha}(x) := e^{-x} x^{n+\alpha}$ is analytic in $\{z \in \mathbb{C} \mid \operatorname{Re} z > 0\}$. By (27) we have

$$x^\alpha e^{-x} L_n^{(\alpha)}(x) = \frac{1}{n!} f_{n\alpha}^{(n)}(x) .$$

If $x \in \mathbb{R}^+$, then for $z = x + re^{i\theta}$ ($r \in (0, x)$) we have by Taylor's formula

$$f_{n\alpha}(z) = \sum_{k=0}^{\infty} \frac{1}{k!} f_{n\alpha}^{(k)}(x) (z-x)^k ,$$

and Cauchy's integral formula gives the estimate

$$\frac{1}{k!} \left| f_{n\alpha}^{(k)}(x) \right| < \frac{1}{r^k} \max_{0 \leq \theta \leq 2\pi} |f_{n\alpha}(x + re^{i\theta})| .$$

Especially for $k = n$ this yields

$$x^\alpha e^{-x} \left| L_n^{(\alpha)}(x) \right| r^n < \max_{0 \leq \theta \leq 2\pi} |f_{n\alpha}(x + re^{i\theta})| \quad (29)$$

If furthermore $n + \alpha > 0$, then $f_{n\alpha}$ is analytic in all of $\mathbb{C}$, and (29) holds for all $r \in \mathbb{R}^+$. This case will be studied now.

To give an estimate of $\max_{0 \leq \theta \leq 2\pi} |f_{n\alpha}(x + re^{i\theta})|$, we expand

$$f_{n\alpha}(x + re^{i\theta}) = e^{-(x+re^{i\theta})} (x + re^{i\theta})^{n+\alpha}$$

to get

$$\max_{0 \leq \theta \leq 2\pi} |f_{n\alpha}(x + re^{i\theta})| = e^{-x} \max_{0 \leq \theta \leq 2\pi} e^{-r \cos \theta} (x^2 + r^2 + 2rx \cos \theta)^{\frac{n+\alpha}{2}} .$$

Together with (29) we have therefore

$$x^\alpha \left| L_n^{(\alpha)}(x) \right| r^n < \max_{0 \leq \theta \leq 2\pi} e^{-r \cos \theta} (x^2 + r^2 + 2rx \cos \theta)^{\frac{n+\alpha}{2}} .$$

We set now $\lambda := \cos \theta \in [-1, 1]$, $p(\lambda) := x^2 + r^2 + 2rx\lambda$, and $q(\lambda) := e^{-r\lambda} p(\lambda)^{\frac{n+\alpha}{2}}$, and have therefore

$$x^\alpha \left| L_n^{(\alpha)}(x) \right| r^n < \max_{-1 \leq \lambda \leq 1} q(\lambda) . \quad (30)$$

As

$$q'(\lambda) = r e^{-r\lambda} p(\lambda)^{\frac{n+\alpha-2}{2}} (x(n+\alpha) - p(\lambda))$$

we get for a possible critical point λ_0 of q the relation

$$\lambda_0 = \frac{x(n+\alpha) - (x^2 + r^2)}{2rx} .$$

At the point $\lambda = \lambda_0$ we have furthermore

$$p(\lambda_0) = x^2 + r^2 + 2rx\lambda_0 = x(n+\alpha) > 0 ,$$

hence

$$q''(\lambda_0) = -2r^2 x e^{-r\lambda_0} p(\lambda_0)^{\frac{n+\alpha-2}{2}} < 0 ,$$

and λ_0 maximizes q . Therefore, from (30) we get

$$x^\alpha \left| L_n^{(\alpha)}(x) \right| r^n < q(\lambda_0) \quad (31)$$

if $-1 \leq \lambda_0 \leq 1$.

We consider now the case $x \in (0, 4(n+\alpha)]$ (which with respect to the representation (13) corresponds to the critical region $(0, 2n)$ for $t = x/2$), and choose $r := \sqrt{x(n+\alpha)}$. In this case we have $\lambda_0 = \frac{1}{2} \sqrt{\frac{x}{n+\alpha}} \in [-1, 0)$. Hence, (31) implies

$$x^\alpha \left| L_n^{(\alpha)}(x) \right| (x(n+\alpha))^{\frac{n}{2}} < e^{\frac{x}{2}} (x(n+\alpha))^{\frac{n+\alpha}{2}} ,$$

and therefore finally

Theorem 2 For the generalized Laguerre polynomials $L_n^{(\alpha)}$ ($\alpha \in \mathbb{Z}$) the estimate

$$e^{-\frac{x}{2}} \left| L_n^{(\alpha)}(x) \right| < \left(\frac{n+\alpha}{x} \right)^{\frac{\alpha}{2}} \quad (32)$$

holds for $x \in (0, 4(n+\alpha)]$ if $n+\alpha > 0$. □

If we define the functions ($\alpha \in \mathbb{R}$)

$$F_n^{(\alpha)}(t) := e^{-t} L_n^{(\alpha)}(2t) \quad (33)$$

then (32) reads ($x = 2t$)

$$|F_n^{(\alpha)}(t)| < \left(\frac{n+\alpha}{2t} \right)^{\frac{\alpha}{2}} \quad (t \in (0, 2(n+\alpha)]) .$$

For $\alpha = 0$, we have Szegő's result (28) in this interval, and for $\alpha = 1$ we get in view of representation (13)

$$|F_n(t)| < \sqrt{\frac{2t}{n}} \quad (t \in (0, 2n)) . \quad (34)$$

This inequality improves (26) as a Puiseux expansion yields

$$\sqrt{\frac{4t}{2n-t}} = \sqrt{\frac{2t}{n}} + \frac{1}{2\sqrt{2}} \left(\frac{t}{n}\right)^{\frac{3}{2}} + P\left(\frac{t}{n}\right)$$

with some positive function P .

Note that the special choice $\alpha = -1$ (and *not* the value $\alpha = 0$) generates the Bateman functions $F_n(t) = F_n^{(-1)}(t)$.

6 Asymptotic estimates

We consider the functions ($\alpha \in \mathbb{R}$)

$$F_n^{(\alpha)}(t) = e^{-t} L_n^{(\alpha)}(2t) = e^{-t} \left(L_n^{(\alpha+1)}(2t) - L_{n-1}^{(\alpha+1)}(2t) \right) = F_n^{(\alpha+1)}(t) - F_{n-1}^{(\alpha+1)}(t) \quad (35)$$

of (33) (see [23], p. 216, formula (1.15)) now in more detail. Taking derivative yields

$$\begin{aligned} \left(F_n^{(\alpha)} \right)'(t) &= -F_n^{(\alpha)}(t) + 2e^{-t} \left(L_n^{(\alpha)} \right)'(2t) \\ &\stackrel{(35)}{=} -e^{-t} \left(L_n^{(\alpha+1)}(2t) - L_{n-1}^{(\alpha+1)}(2t) \right) + 2e^{-t} \left(L_n^{(\alpha)} \right)'(2t) \\ &= -e^{-t} \left(L_n^{(\alpha+1)}(2t) + L_{n-1}^{(\alpha+1)}(2t) \right) \\ &= - \left(F_n^{(\alpha+1)}(t) + F_{n-1}^{(\alpha+1)}(t) \right) . \end{aligned} \quad (36)$$

where the relation about $\left(L_n^{(\alpha)} \right)'$ corresponds to ([23], p. 215, formula (1.12)).

Moreover the program [12] generates the differential equation

$$(1 + \alpha + 2n - t) F_n^{(\alpha)}(t) + (1 + \alpha) \left(F_n^{(\alpha)} \right)'(t) + t \left(F_n^{(\alpha)} \right)''(t) = 0$$

for the functions $F_n^{(\alpha)}$ with respect to the variable t , and the recurrence equation

$$(-1 + \alpha + n) F_{n-2}^{(\alpha)} + (1 - \alpha - 2n + 2t) F_{n-1}^{(\alpha)} + n F_n^{(\alpha)} = 0$$

with respect to the variable n (check!).

Assume now, $t_1 < t_2$. Then we get by an integration

$$\begin{aligned} \left(F_n^{(\alpha)} \right)^2(t_2) - \left(F_n^{(\alpha)} \right)^2(t_1) &= 2 \int_{t_1}^{t_2} F_n^{(\alpha)}(t) \left(F_n^{(\alpha)} \right)'(t) dt \\ &\stackrel{(35),(36)}{=} -2 \int_{t_1}^{t_2} \left(\left(F_n^{(\alpha+1)}(t) \right)^2 - \left(F_{n-1}^{(\alpha+1)}(t) \right)^2 \right) dt . \end{aligned}$$

If we choose $\alpha = 0$, we get in particular

$$\left(F_n^{(0)}\right)^2(t_2) - \left(F_n^{(0)}\right)^2(t_1) = -2 \int_{t_1}^{t_2} \left(\left(F_n^{(1)}(t)\right)^2 - \left(F_{n-1}^{(1)}(t)\right)^2 \right) dt. \quad (37)$$

Together with the relation ($n \in \mathbb{N}$)

$$F_n(t) = F_n^{(-1)}(t) = -\frac{2t}{n} F_{n-1}^{(1)}(t)$$

(see (12) and (13), or [22], p. 98, formula (5.2.1)) it followffs from (37) that

$$\left(F_n^{(0)}\right)^2(t_2) - \left(F_n^{(0)}\right)^2(t_1) = \int_{t_1}^{t_2} F_{n+1}^2(t) \frac{(n+1)^2}{2t^2} dt - \int_{t_1}^{t_2} F_n^2(t) \frac{n^2}{2t^2} dt.$$

We now let $t_2 \rightarrow \infty$. Then $F_n^{(0)}(t_2) \rightarrow 0$, and as

$$\int_0^\infty F_n^2(t) dt = 1 \quad (38)$$

(see (19), compare [3], formulae (2.7) and (2.91)), we get further

$$\begin{aligned} \left(F_n^{(0)}\right)^2(t_1) &= \int_{t_1}^\infty F_{n+1}^2(t) \frac{(n+1)^2}{2t^2} dt - \int_{t_1}^\infty F_n^2(t) \frac{n^2}{2t^2} dt \\ &\leq \int_{t_1}^\infty F_{n+1}^2(t) \frac{(n+1)^2}{2t^2} dt \leq \frac{(n+1)^2}{2t_1^2} \left(1 - \int_0^{t_1} F_{n+1}^2(t) dt\right) \end{aligned}$$

where the last inequality is deduced with (38). Soff we have finally the inequality ($n \in \mathbb{N}$, $t > 0$)

$$\left|F_n^{(0)}(t)\right| \leq \frac{n+1}{\sqrt{2}t}.$$

This sharpens the result of Szegö (28) for large t . From (35) and (36) it follows that

$$F_n^{(\alpha)}(t) + \left(F_n^{(\alpha)}\right)'(t) = -2 F_{n-1}^{(\alpha+1)}(t).$$

We deduce now for a critical point t_k of F_n with $F_n'(t_k) = 0$ the relation

$$|F_n(t_k)| = 2 \left|F_{n-1}^{(0)}(t_k)\right| \leq \sqrt{2} \frac{n}{t_k}. \quad (39)$$

Especially is this $\leq 2/e$ for

$$\frac{t_k}{n} \geq \frac{e}{\sqrt{2}} \approx 1.92211551407955841\dots \quad (40)$$

It is now remarkable that by the result of Hahn (21) for $n \rightarrow \infty$ the most important critical point T of F_n which produces the maximal value of F_n has the property $T/n \rightarrow 2$ as $T_n < T < 2$. This gives finally the following

Theorem 3 The Krzyż conjecture is asymptotically true for the superordinate functions $e^{-t \frac{1+z}{1-z}}$, i.e. we have $\left| a_n \left(e^{-t \frac{1+z}{1-z}} \right) \right| \leq \frac{2}{e}$ for $n \geq N$. $\square$

We will now strengthen this result.

Therefore let an arbitrary positive zero t_n of F_n be given. Then t_n is also a zero of $h_n := F_n^2$, and as

$$h_n''(t) = 2 (F_n'(t))^2 + 2 F_n(t) F_n''(t) ,$$

by the differential equation for F_n we get

$$h_n''(t) = 2 (F_n'(t))^2 + 2 \frac{t-2n}{t} F_n^2(t) . \quad (41)$$

From this we may deduce that $h_n''(t) > 0$ for $t \geq 2n$. Now, however, we consider the interval between t_n and the smallest relative extremum $t_n^* > t_n$ of F_n , i. e. the smallest zero t_n^* of F_n' after t_n . Then obviously h_n is strictly increasing in $[t_n, t_n^*]$, further $h_n'(t_n) = h_n'(t_n^*) = 0$, and therefore $h_n' \Big|_{[t_n, t_n^*]}$ assumes an absolute maximum at some interior point $t_n^{**} \in (t_n, t_n^*)$, where $h_n''(t_n^{**}) = 0$.

From (41) we deduce

$$(F_n'(t_n^{**}))^2 = \frac{2n - t_n^{**}}{t_n^{**}} F_n^2(t_n^{**}) ,$$

and therefore by (18)

$$|F_n'(t_n^{**})| = \sqrt{\frac{2n - t_n^{**}}{t_n^{**}}} |F_n(t_n^{**})| < \sqrt{\frac{2n - t_n^{**}}{t_n^{**}}} .$$

As $\sqrt{\frac{2n-t}{t}}$ is strictly decreasing for $t \in (0, 2n)$, it follows furthermore that

$$|F_n'(t_n^{**})| < \sqrt{\frac{2n - t_n}{t_n}} ,$$

and finally (h_n' is positive)

$$h_n'(t_n^{**}) = 2 |F_n(t_n^{**})| |F_n'(t_n^{**})| < 2 \sqrt{\frac{2n - t_n}{t_n}}$$

using (18) again. As t_n^{**} is the global maximum of h_n' in $[t_n, t_n^*]$, we therefore are lead to the inequalities

$$0 < h_n'(t) < 2 \sqrt{\frac{2n - t_n}{t_n}} \quad (42)$$

for all $t \in (t_n, t_n^*)$.

We are interested in $h_n(t_n^*)$, the value of h_n at its maximum t_n^* . Therefore let $p > 0$ be given such that $h_n(t_n^*) > \frac{1}{p}$. As $h_n(t_n) = 0$, and h_n is stricly increasing, there is some $\tilde{t}_n \in (t_n, t_n^*)$ with $h_n(\tilde{t}_n) = \frac{1}{p}$. The mean value theorem then shows the existence of $\xi_n \in (\tilde{t}_n, t_n^*)$ with

$$\frac{h_n(t_n^*) - h_n(\tilde{t}_n)}{t_n^* - \tilde{t}_n} = h_n'(\xi_n) ,$$

and therefore by (42)

$$\frac{h_n(t_n^*) - h_n(\tilde{t}_n)}{t_n^* - \tilde{t}_n} < 2 \sqrt{\frac{2n - t_n}{t_n}}$$

or

$$h_n(t_n^*) < h_n(\tilde{t}_n) + 2 \sqrt{\frac{2n - t_n}{t_n}} (t_n^* - \tilde{t}_n) = \frac{1}{p} + 2 \sqrt{\frac{2n - t_n}{t_n}} (t_n^* - \tilde{t}_n).$$

By (19) we have

$$\int_0^\infty h_n(\tau) d\tau = 1,$$

and thus by the integral mean value theorem (h_n is nonnegative)

$$1 > \int_{\tilde{t}_n}^{t_n^*} h_n(\tau) d\tau = h_n(\eta_n) (t_n^* - \tilde{t}_n)$$

for some $\eta_n \in (\tilde{t}_n, t_n^*)$. As h_n is strictly increasing, we therefore get $h_n(\eta_n) > h_n(\tilde{t}_n) = \frac{1}{p}$ implying

$$1 > \frac{1}{p} (t_n^* - \tilde{t}_n) \quad \text{or} \quad t_n^* - \tilde{t}_n < p.$$

Finally we have

$$h_n(t_n^*) < \frac{1}{p} + 2p \sqrt{\frac{2n - t_n}{t_n}}.$$

We were lead to this inequality under the assumption that $h_n(t_n^*) > \frac{1}{p}$. If $h_n(t_n^*) \leq \frac{1}{p}$, however, then the same conclusion follows trivially, so that the above calculations can be summarized by the following

Lemma 1 Let $h_n(t) = F_n^2(t)$, let t_n be a positive zero of F_n , let t_n^* the lowest zero of F_n' that is larger than t_n , and let $p > 0$. Then

$$h_n(t_n^*) < \frac{1}{p} + 2p \sqrt{\frac{2n - t_n}{t_n}}. \quad \square$$

We now emphasize on the largest zero $t_n = T_n$ of F_n . By the results of § 2 the global maximum of F_n is attained at the last zero T_n^* of F_n' which lies in the interval $(T_n, 2n)$, and is therefore the smallest zero of F_n' after T_n . So Lemma 1 applies.

By a result of Bottema and Hahn (see [5], anffd [8], p. 228, last formula), the inequalityff

$$T_n > 2n - \frac{3}{2} - 8\sqrt{2}\sqrt{n-1} =: \tau_n \quad (43)$$

($n \geq 33$) holds for the last zero of F_n (or $L_{n-1}^{(1)}$). As $\sqrt{\frac{2n-t}{t}}$ is strictly decreasing for $t \in (0, 2n)$, we have the inequality

$$\sqrt{\frac{2n - T_n}{T_n}} < \sqrt{\frac{2n - \tau_n}{\tau_n}}.$$

Puiseux expansion yields the asymptotic expression ($n \rightarrow \infty$)

$$\sqrt{\frac{2n - \tau_n}{\tau_n}} = 2 \sqrt[4]{2} \frac{1}{n^{1/4}} + \frac{131}{16 \sqrt[4]{2}} \frac{1}{n^{3/4}} + O\left(\frac{1}{n^{5/4}}\right),$$

especially is

$$\sqrt{\frac{2n - \tau_n}{\tau_n}} \sim \frac{1}{n^{1/4}}.$$

In our calculations the value p was arbitrary, so we have the freedom to choose it properly. The asymptotics suggest the choice $p \sim n^{1/8}$. For any $a > 0$ we get therefore

$$h_n(T_n^*) < \frac{a}{n^{1/8}} + 2 \frac{n^{1/8}}{a} \sqrt{\frac{2n - T_n}{T_n}} < \frac{a}{n^{1/8}} + 2 \frac{n^{1/8}}{a} \sqrt{\frac{2n - \tau_n}{\tau_n}} \sim \frac{1}{n^{1/8}} < \frac{b}{n^{1/8}}$$

for some $b > 0$.

We choose the value $a = 2 \sqrt[8]{2}$ (minimizing the leading term in the corresponding Puiseux expansion) and get the global estimate

$$h_n(T_n^*) < 2 \sqrt[8]{2} \frac{1}{n^{1/8}} + \frac{\sqrt{\frac{3}{2} + 8 \sqrt{2} \sqrt{n-1}}}{\sqrt[8]{2} \sqrt{2n - \frac{3}{2} - 8 \sqrt{2} \sqrt{n-1}}} n^{1/8}.$$

Now we remember that F_n takes its global maximum over $\mathbb{R}^+$ at the point T_n^* , and so does h_n . We therefore have for all $a > 0$, $n \in \mathbb{N}$ and $t > 0$ the inequality

$$|F_n(t)| < \sqrt{2 \sqrt[8]{2} \frac{1}{n^{1/8}} + \frac{\sqrt{\frac{3}{2} + 8 \sqrt{2} \sqrt{n-1}}}{\sqrt[8]{2} \sqrt{2n - \frac{3}{2} - 8 \sqrt{2} \sqrt{n-1}}} n^{1/8}} = \frac{2 \sqrt[16]{2}}{n^{1/16}} + \frac{131}{64 \cdot 2^{7/16} n^{9/16}} + O\left(\frac{1}{n^{17/16}}\right). \quad (44)$$

We mention that we get a better asymptotic estimate if we use the sharper left hand inequality (20) instead of (43), set $\tau_n^* := -1 + 2n - C(2n - 1)$ (C constant) leading to the asymptotic result

$$\sqrt{\frac{2n - \tau_n^*}{\tau_n^*}} \sim \frac{1}{n^{1/3}},$$

and therefore by the choice $p \sim n^{1/6}$ and the same procedure as above to the

Theorem 4 For all $t \in \mathbb{R}^+$ we have the asymptotic inequality ($n \geq N$)

$$|F_n(t)| < \frac{c}{n^{1/12}}$$

for some $c > 0$, and in particular the limiting value

$$\lim_{n \rightarrow \infty} |F_n(t)| = 0.$$

□

Obviously this theorem strengthens Theorem 3.

In principle (44) enables one to prove the statement

$$|F_n(t)| \leq \frac{2}{e}$$

for all $n \in \mathbb{N}$. Therefore one shows that the estimation function

$$E(n) = \sqrt{2 \sqrt[8]{2} \frac{1}{n^{1/8}} + \frac{\sqrt{\frac{3}{2} + 8\sqrt{2}\sqrt{n-1}}}{\sqrt[8]{2}\sqrt{2n - \frac{3}{2} - 8\sqrt{2}\sqrt{n-1}}}} n^{1/8}$$

of (44) is decreasing, and as $E(17821075) > 2/e$ and $E(17821076) < 2/e$, it remains to prove the result for only a finite number of initial values.

The number of initial values, however, can be decisively fflowered using that by (40) $|F_n(t)| \leq 2/e$ whenever $T_n^*/n \geq \frac{e}{\sqrt{2}}$, especially if $T_n/n \geq \frac{e}{\sqrt{2}}$. From the Bottema-Hahn bound

$$\frac{T_n}{n} > \frac{\tau_n}{n} = 2 - \frac{3}{2n} - 8\sqrt{2} \frac{\sqrt{n-1}}{n} =: e(n)$$

we obtain first by the calculation

$$e'(n) = \frac{-16\sqrt{2} + 3\sqrt{n-1} + 8\sqrt{2}n}{2n^2\sqrt{n-1}}$$

that $e(n)$ is increasing for $n \geq 2$, and as $\lim_{n \rightarrow \infty} e(n) = 2$ there is exactly one solution $n_0 \geq 2$ of the equation $e(n) = \frac{e}{\sqrt{2}}$, and $T_n/n > e(n) \geq e(n_0) = \frac{e}{\sqrt{2}}$ for $n > n_0$. A numerical calculation shows that $n_0 \approx 21138.7$ so that we are lead to the

Theorem 5 The inequality $|F_n(t)| \leq 2/e$ is true for all $n \in \mathbb{N}$ and all $t > 0$ if it is true for $n \leq 21138$.

7 Estimates for the derivative

By (23) it follows that at the zeros of F_n the derivative F'_n satisfies the relation $|F'_n(t)| \leq 2$. This result holds for all $t \geq 0$ which can be seen as follows: Using (36) with $\alpha = -1$ we have

$$F'_n(t) = -e^{-t} \left(L_n(2t) + L_{n-1}(2t) \right), \quad (45)$$

and by an application of the Szegö result (28) it follows for $t > 0$

$$|F'_n(t)| < 2. \quad (46)$$

This shows that for all $n \in \mathbb{N}$ the derivatives $|F'_n|$ have their maximal value at the origin where $|F'_n(0)| = 2$ (see (11)). We note moreover, even though ffthe derivative $F := F_n$ has a representation (45) similar to that of F_n itself, it satisfies the much more complicated differential equation

$$\left(-4n^2 + 4nt - t^2 \right) F(t) - 2nF'(t) + \left(-2nt + t^2 \right) F''(t) = 0.$$

8 The functions H_n

In this section we collect the explicit inequalities that we deduced, and formulate a conjecture concerning the Bateman functions.

As the last point of inflection of the functions F_n is at the point $t = 2n$ which increases with increasing n , it is reasonable to introduce the functions

$$H_n(t) := (-1)^n F_n(nt)$$

that have common absolute values with F_n which, however, are attained at different points. The scale on the t -axis is here such that the last point of inflection lies at $t = 2$ for all functions H_n ($n \in \mathbb{N}$), and H_n is positive for $t \geq 2$. It is easy to deduce the differential equation

$$t H_n''(t) = n^2 (t - 2) H_n(t) \quad (47)$$

satisfied by H_n .




Figure 3: Estimates for the functions H_n

The inequalities that we deduced for F_n read as follows for H_n : The trivial estimate (18) gives

$$|H_n(t)| \leq 1 ,$$

(16) yields ($n > 2$)

$$|H_n(t)| \leq \frac{2}{t} ,$$

the refinement (39) gives

$$|H_n(t_k)| \leq \frac{\sqrt{2}}{t_k}$$

for a critical point t_k of H_n , (26) implies

$$|H_n(t)| < \sqrt{\frac{4t}{2-t}} ,$$

and finally (34) yields

$$|H_n(t)| < \sqrt{2t}.$$

These estimates commonly do not depend on n . One more estimate will be added in the next section. Figure 3 shows them graphically.

Figures 4 and 5 show the graphs of the functions H_n ($n = 1, \dots, 20$). We conjecture that H_n is strictly decreasing for increasing n at the point $t = 2$. Note that by the result of Hahn (21) this is not true for any $t < 2$. In the next section we will show, however, that $\lim_{n \rightarrow \infty} H_n(t) = 0$ for each $t > 2$.




Figure 4: The functions H_n ($n = 1, \dots, 20$)




Figure 5: The functions H_n ($n = 1, \dots, 20$) in the interval $[1.5, 2]$

9 Estimates for large t

In this section we show that for all $t > 2$ the values $H_n(t)$ tend to 0 for $n \rightarrow \infty$.

The inequalities (46), and (17) correspond to the inequalities

$$|H'_n(t)| \leq 2n ,$$

and

$$|H'_n(t)| \leq \frac{n}{t} \quad (48)$$

for the derivative of H_n , respectively.

By the differential equation (47) for H_n we have

$$H_n(t) = \frac{t}{n^2(t-2)} H''_n(t) . \quad (49)$$

Let now $2 < t_1 < t_2$ be given. By definition H_n is strictly positive in $[t_1, t_2]$, and by (49) H''_n is strictly positive in $[t_1, t_2]$. Therefore

$$\int_{t_1}^{t_2} H_n(t) dt = \frac{1}{n^2} \int_{t_1}^{t_2} \frac{t}{t-2} H''_n(t) dt .$$

As the function $t/(t-2)$ is strictly decreasing, we have

$$\max_{t \in [t_1, t_2]} \frac{t}{t-2} = \frac{t_1}{t_1-2} ,$$

and therefore

$$\frac{1}{n^2} \int_{t_1}^{t_2} \frac{t}{t-2} H''_n(t) dt \leq \frac{1}{n^2} \frac{t_1}{t_1-2} \int_{t_1}^{t_2} H''_n(t) dt .$$

As H''_n is positive in $[t_1, t_2]$, and as $\lim_{t \rightarrow \infty} H'_n(t) = 0$, it follows that H'_n is negative and increasing, and therefore

$$\int_{t_1}^{t_2} H_n(t) dt \leq \frac{1}{n^2} \frac{t_1}{t_1-2} |H'_n(t_1) - H'_n(t_2)| \leq \frac{1}{n^2} \frac{t_1}{t_1-2} |H'_n(t_1)| \leq \frac{1}{n^2} \frac{t_1}{t_1-2} \frac{n}{t_1} \leq \frac{1}{n(t_1-2)}$$

where we used (48). For fixed $t_2 > 2$ we set now $t_1 := \frac{2+t_2}{2}$, and with the integral mean value theorem we find $\tau \in [t_1, t_2]$ with

$$H(t_2) \leq H(\tau) = \frac{1}{t_2 - t_1} \int_{t_1}^{t_2} H_n(t) dt \leq \frac{1}{(t_2 - t_1)(t_1 - 2)} \frac{1}{n} = \frac{4}{(t_2 - 2)^2} \frac{1}{n} .$$

Another estimate for large t which is independent of n , will be established now. Let again $2 < t_1 < t_2$. Then by (49)

$$\begin{aligned} |H_n^2(t_2) - H_n^2(t_1)| &= \left| \int_{t_1}^{t_2} (H_n^2(t))' dt \right| = \left| \int_{t_1}^{t_2} 2 H'_n(t) H_n(t) dt \right| = \left| \int_{t_1}^{t_2} \frac{t}{n^2(t-2)} H'_n(t) H''_n(t) dt \right| \\ &\leq \frac{t_1}{n^2(t_1-2)} \int_{t_1}^{t_2} |2 H'_n(t) H''_n(t)| dt = \frac{t_1}{n^2(t_1-2)} |(H'_n)^2(t_2) - (H'_n)^2(t_1)| . \end{aligned}$$

We let now $t_2 \rightarrow \infty$, and get ($t := t_1$) using (48)

$$|H_n(t)|^2 \leq \frac{t}{n^2(t-2)} |(H'_n)^2(t)| \leq \frac{1}{t(t-2)}.$$

This estimate improves the earlier ones for large t , see Figure 3.

Finally we show that uniformly with respect to n the functions H_n decrease faster than each negative power.

Theorem 6 For each $k \in \mathbb{N}_0$ there is a constant $C_k > 0$ such that

$$|H_n(t)| \leq \frac{C_k}{t^k} \quad (t > 0, n \in \mathbb{N})$$

that is independent of n .

Proof: We prove the result by induction with respect to k . For $k = 0$ the statement is trivially true, see (18). Assume now the statement holds for some $k \in \mathbb{N}_0$, i. e.

$$|F_n(t)| \leq C_k \left(\frac{n}{t}\right)^k.$$

Then we get using (14)

$$\begin{aligned} |F_n(t)| &= \frac{1}{2t} |(n-1)(F_n(t) - F_{n-1}(t)) + (n+1)(F_n(t) - F_{n+1}(t))| \\ &\leq \frac{2(n-1)C_k + (n+1)C_k + (n+1)C_k D_k}{2t} \left(\frac{n}{t}\right)^k \\ &\leq \frac{2(n-1)C_k D_k + (n+1)C_k D_k + (n+1)C_k D_k}{2t} \left(\frac{n}{t}\right)^k \\ &= 2C_k D_k \left(\frac{n}{t}\right)^{k+1}, \end{aligned}$$

where we chose $D_k \geq 1$ such that $(n+1)^k \leq D_k n^k$ (the choice $D_k = 2^k$ does the job required as $(n+1)^k \leq (2n)^k \leq 2^k n^k$). This yields the result. $\square$

References

- [1] Abramowitz, M. and Stegun, I. A.: *Handbook of Mathematical Functions*. Dover Publ., New York, 1964.
- [2] Amer. Math. Monthly **59** (1952), 45, Problem 4468, and solution in Amer. Math. Monthly **60** (1953), 131–132.
- [3] Bateman, H.: The k -function, a particular case of the confluent hypergeometric function. Trans. Amer. Math. Soc. **33** (1931), 817–831.
- [4] Behnke, H. and Sommer, F.: *Theorie der analytischen Funktionen einer komplexen Veränderlichen*. Grundlehren der mathematischen Wissenschaften **77**, Springer, Berlin–Heidelberg–New York, 1965.

- [5] Bottema, O.: Die Nullstellen gewisser durch Rekursionsformeln definierter Polynome. Proc. Amsterdam **34** (1930), 681–691.
- [6] Doetsch, G.: *Einführung in Theorie und Anwendung der Laplace-Transformation*. Mathematische Reihe **24**, Birkhäuser, Basel–Stuttgart, 1958.
- [7] Duren, P. L.: *Univalent functions*. Grundlehren der mathematischen Wissenschaften **259**, Springer-Verlag, New York–Berlin–Heidelberg–Tokyo, 1983.
- [8] Hahn, W.: Bericht über die Nullstellen der Laguerreschen und der Hermiteschen Polynome. Jber. DMV **44** (1934), 215–236.
- [9] Hoffman, K.: Banach spaces of analytic functions. Prentice-Hall Series in Modern Analysis, Prentice-Hall, Englewood Cliffs, N. J., 1996.
- [10] Hummel, J. A., Scheinberg, St. and Zalcman, L.: A coefficient problem for bounded nonvanishing functions. J. Anal. Math. **31** (1977), 169–190.
- [11] Koepf, W.: Power series in Computer Algebra. J. Symb. Comp. **13** (1992) 581–603.
- [12] Koepf, W.: MATHEMATICA package `PowerSeries.m`. Fachbereich Mathematik, Freie Universität Berlin, 1991–1993.
- [13] Krzyż, J.: Problem 1, posed in: Fourth Conference on Analytic Functions, Ann. Polon. Math. **20** (1967–1968), 314.
- [14] Levin, V.: Aufgabe 163, Jber. DMV **43** (1934), 113.
- [15] Littlewood, J. E.: *Lectures on the theory of functions*. Oxford University Press, London, 1944.
- [16] Rajagopal, C.T.: On inequalities for analytic functions. Amer. Math. Monthly **60** (1953), 693–695.
- [17] Reissner, E.: Lösung der Aufgabe 163, Jber. DMV **44** (1934), 80–83.
- [18] Robertson, M. S.: Solution of Problem 4468, Amer. Math. Monthly **60** (1953), 131–132.
- [19] Ryff, J. V.: Subordinate H^p functions.
- [20] Shapiro, H. S.: Problem 4468, Amer. Math. Monthly **59** (1952), 45.
- [21] Szegő, G.: Ein Beitrag zur Theorie der Polynome von Laguerre und Jacobi. Math. Z. **1** (1918), 341–356.
- [22] Szegő, G.: *Orthogonal Polynomials*. Amer. Math. Soc. Coll. Publ. Vol. **23**, New York City, 1939.
- [23] Tricomi, F. G.: *Vorlesungen über Orthogonalreihen*. Grundlehren der Mathematischen Wissenschaften **76**, Springer-Verlag, Berlin–Göttingen–Heidelberg, 1955.